

POLYCYCLIC AROMATIC HYDROCARBON BIODEGRADATION AS A FUNCTION OF OXYGEN TENSION IN CONTAMINATED SOIL

C.J. Hurst¹, R.C. Sims¹, J.L. Sims¹, D.L. Sorensen¹, J.E. McLean¹ and S. Huling², ¹Division of Environmental Engineering, Department of Civil and Environmental Engineering, and the Utah Water Research Laboratory, Utah State University, Logan, UT 84322-8200 and ²U.S. EPA, Robert S. Kerr Environmental Research Laboratory, Ada, OK

ABSTRACT

Laboratory tests were conducted to determine the effect of soil gas oxygen concentration on the degradation and mineralization of spiked ¹⁴C-pyrene and nonspiked 16 priority pollutant polycyclic aromatic hydrocarbons (PAHs) present in the soil. The soil used for the evaluation was taken from a prepared bed land treatment unit at the Champion International Superfund Site in Libby, Montana. This soil was contaminated with wood preserving wastes including creosote composed primarily of polycyclic aromatic hydrocarbons and pentachlorophenol. Degradation rates of ¹⁴C-pyrene and PAH compounds were found to be enhanced under soil gas oxygen concentrations between 2% and 21% in the contaminated soil. Between 45% and 55% of ¹⁴C-pyrene spiked onto the soil was mineralized after 70 days at soil gas oxygen levels between 2% and 21%. No statistically significant mineralization was found to occur at 0% oxygen concentrations. Mineralization of ¹⁴C-pyrene in contaminated soil poisoned with mercuric chloride was determined to be less than 0.5%. Degradation of indigenous nonradiolabeled PAHs in nonpoisoned soil was statistically significantly greater than in poisoned soil. These results indicated that degradation of ¹⁴C-pyrene and PAH compounds was biological and would occur under low oxygen concentrations that could be maintained, for example, using bioventing technology in order to achieve continued treatment of buried lifts of soil while new lifts are added, thus decreasing the total time for soil remediation of the prepared bed.

KEY WORDS

bioremediation, prepared-bed bioreactor, oxygen tension, contaminated soil

INTRODUCTION

Polycyclic aromatic hydrocarbons (PAHs) include a group of organic priority pollutants of critical environmental and public health concern due to the following characteristics: (1) chronic health effects (carcinogenicity); (2) microbial recalcitrance; (3) high bioaccumulation potential; and (4) low removal efficiencies in traditional treatment processes [1]. PAHs are classified as both carcinogenic and non-carcinogenic compounds. Carcinogenic PAHs refer to fluoranthene, pyrene, benzo(a)anthracene, chrysene, benzo(b)fluoranthene, benzo(k)fluoran-

thene, benzo(a)pyrene, dibenzo(a,h)anthracene, benzo(g,h,i)perylene and indeno(1,2,3-cd)pyrene. Non-carcinogenic PAHs refer to naphthalene, acenaphthylene, acenaphthene, fluorene, anthracene and phenanthrene. These compounds are produced by high-temperature industrial processes such as petroleum refining, coke production, wood preservation, and synthetic oil and gas production. PAHs are associated with a wide range of hazardous waste sites (e.g., Superfund sites, uncontrolled hazardous waste sites not listed as Superfund sites, and hazardous waste land treatment sites) [2, 3].

Though PAHs are considered recalcitrant, losses do occur over time. These losses may occur through biotic processes and a variety of abiotic processes including leaching, photodegradation and volatilization [4]. Several environmental factors are known to influence the capacity of indigenous microbial populations to degrade PAHs. The interactions among environmental factors such as temperature, pH, soil gas oxygen concentrations, oxidation-reduction potential and the presence of other substrates often control the feasibility of biodegradation [3, 5, 6]. However, the soil gas oxygen concentration is generally thought to be the limiting constituent in the biodegradation of PAHs when nutrients are not limiting [7].

In order to achieve biodegradation of PAH compounds, these electron donors must be brought into contact with electron acceptors and with metabolically active microorganisms. It has been shown that the initial attack on the reduced hydrocarbon PAH molecule is an aerobic one and involves oxygen [8-10]. Dihydroxylation of the aromatic nucleus is a prerequisite for enzymatic cleavage of the benzene ring by oxygenases that incorporates molecular oxygen into the ring [11]. The incorporation of molecular oxygen into the PAH molecule is an activation step that makes the molecule biologically active and also a substrate for ring fission. A demonstration that molecular oxygen is involved in hydrocarbon utilization by microorganisms was presented by Hansen and Kallio [12]. Consequently, molecular oxygen is required for the insertion of oxygen into the hydrocarbon molecule, but not for the subsequent utilization of the oxygenated intermediates. Park and Sims [7] demonstrated that the concentration of oxygen was strongly correlated with the rate of apparent disappearance of 7,12-dimethylbenzanthracene spiked onto unsaturated soil in laboratory flask reactors.

In the presence of the appropriate organisms, biodegradation of lower molecular

weight PAHs such as naphthalene occurs rapidly. Biodegradation of 3-ring PAH compounds such as acenaphthene, anthracene and phenanthrene is much slower. PAH compounds with four or more fused benzene rings are believed to be biodegraded throughout the process of cooxidation, whereby these non-growth PAHs are oxidized when present as cosubstrates in a medium where one or more hydrocarbons are available for growth [13]. Low concentrations of oxygen may limit the cooxidative process as discussed by Sims and Overcash [3]. The initial attack through cooxidation on a recalcitrant molecule such as a PAH is a coincidental attack on that compound that is probably of little consequence to the microorganism involved in the reaction. Neither energy or carbon for biosynthesis results from this oxidation, and the reaction probably results in a loss of energy for the organism involved [14, 15].

Bioremediation of PAHs at field scale is generally accomplished using three types of systems: (1) in situ; (2) prepared bed; or (3) bioreactor (e.g. slurry reactors) systems. This discussion will focus on the biodegradation of PAH in prepared bed systems [16]. This type of contaminated soil treatment generally consists of applying soil layers or lifts of about 6-12 inches into a prepared bed bioreactor also called a Land Treatment Unit (LTU). The LTU is lined with clay and/or plastic liners. A leachate collection system, rainfall runoff/leachate storage, and a passive moisture control system are all components that may be used in LTU design. When the contaminated soil lift reaches the cleanup levels prescribed for the site, a new lift is applied, and the remediation process is continued until all the soil reaches cleanup levels. The LTU is then capped. For microorganisms to get both air and water, the soil moisture content in the LTU is maintained at between 70% and 80% field capacity. Nutrients are also added as needed to the LTU either as solids or dissolved in the irrigation water. Tilling of the top lift is typically conducted

on a weekly basis when weather permits [17, 18].

Subsurface unsaturated soil in prepared-bed systems (buried lifts) is expected to be lower in oxygen concentration with increasing depth through the prepared-bed. This is expected due to a rate of oxygen utilization by microorganisms that is higher than the rate of supply of oxygen through the prepared-bed due to diffusion from the atmosphere. Diffusion has been considered the primary mechanism of oxygen transport in soils [19], and it is known that the oxygen diffusion rate decreases with an increase in soil depth [20].

Because aerobic biological oxidation was found by Lin [21] to be the prime mechanism for the biodegradation of selected compounds of wood-preserving wastes, and because the rate of oxygen diffusion is known to decrease with an increase in soil depth, it is likely that the rate of biodegradation in buried lifts of a prepared-bed system is limited by the anticipated low concentrations of molecular oxygen.

There is currently a lack of information concerning the fate of pollutants in soil at field scale, most importantly those that are considered hazardous to human health and the environment. A small number of cases of land treatment of hazardous wastes have been documented with respect to predicting degradation, immobilization and detoxification of hazardous wastes when mixed with soil and subjected to engineering management [22].


An important problem related to prepared bed bioremediation of wood preservative contaminated unsaturated soil is the length of time required for bioremediation. Treatment time may be shortened by placing a new contaminated soil lift on top of a lower lift before the soil comprising the lower lift reaches target remediation levels. If critical oxygen levels necessary for biodegradation of PAHs are present, then continued bio-

degradation of target chemicals in the lower lift may allow the lower lift to reach target remediation levels as the upper lift begins the bioremediation process.

The soil at the Champion International Superfund Site in Libby, Montana, was contaminated as a result of wood preserving operations between 1946 and 1969. The contaminants mainly consisted of residuals from creosote, primarily PAHs, and pentachlorophenol (PCP) wood preservatives. The prepared bed system at this site consists of two one acre LTUs lined with a 60 mil high density polyethylene liner, compacted clay, and geotextile filter fabric. Nutrients, including ammonium sulfate and ammonium phosphate, are periodically added to the prepared-bed to prevent nutrient limitations to biodegradation. Design criteria for this site were total containment of contaminated soils, surface runoff and leachate, with treatment and disposal of all contaminated soils within the LTU [17, 23].

Oxygen concentrations previously measured in the prepared bed at the site ranged from 0% to 21% in the soil gas phase. Results of sampling at greater depths than the top nine-inch layer (lift) of soil indicated a decrease in oxygen concentration with depth. This decrease is most likely due to the oxygen at the greater depths being consumed rapidly relative to the rate of diffusion from the soil surface [24, 25].

The objective of this study was to evaluate the effect of oxygen concentration in the soil gas phase on the rate and extent of biodegradation of PAHs that occur in wood preservative-contaminated soil undergoing bioremediation within the prepared bed treatment system at the Champion International Superfund Site in Libby, Montana. Results from this research will be used to determine the potential for decreasing the required time for treatment of contaminated soil in prepared bed systems by maintaining the necessary concentration of oxygen in the soil gas phase for active aerobic bio-



degradation of PAHs in buried lifts within the prepared bed.

MATERIALS AND METHODS

Soil Characteristics	Values
Physical	
% sand	48
% silt	39
% clay	13
Texture	Loam
Moisture retention (% water)	
1/3 Bar	12.9
15 Bar	5.5
Chemical	
pH	6.6
EC (mhos/cm)	4.5
Organic Carbon (%)	1.88
CEC (meq/100 g)	6.1
NO ₃ -N (mg/kg)	6.8
Total N (%)	0.10
P (mg/kg)	20.8

Table 1. Soil physical and chemical characteristics.

To determine the effect of oxygen concentration in the soil gas phase on biodegradation in the contaminated soil, the rates of biodegradation of radiolabeled pyrene and the 16 priority pollutant PAH compounds were measured in contaminated soil taken from the prepared bed bioreactor at the Libby site.

Soil characterization

The contaminated soil evaluated in this study was taken from the Prepared-Bed LTU Number One at the Champion International Superfund Site in Libby, Montana [23]. A soil characterization was conducted on the soil taken from the site by the Utah State University Soils Analysis Laboratory. Results are presented in Table 1. The soil was taken from the top six inches of the LTU which had not yet reached the cleanup levels of 88 mg/kg total carcinogenic PAH and 37 mg/kg PCP. Soil moisture for the laboratory study was maintained at 70%-80% of field capacity, the same value maintained at the site.

Mass balance studies of ¹⁴C-pyrene degradation in laboratory microcosms

Radiolabeled pyrene was chosen because it was identified as the rate limiting constituent at the atmospheric oxygen level in soil bioremediation at the site [23]. In addition, the concentrations of unlabeled carcinogenic PAH compounds present in the contaminated soil were measured. A schematic of the glass laboratory biometer flask test reactors that were used for evaluation of the effect of oxygen concentration on PAH biodegradation is shown in Figure 1.

The experiment consisted of evaluating the soil in laboratory biometer flask tests to determine the biodegradation response at five soil gas oxygen concentrations: (1) 0% (anaerobic), (2) 2% O₂, (3) 5% O₂, (4) 10% O₂ and (5) 21% O₂ (atmospheric). Labora-

tory tests were conducted by adding 50 grams of contaminated soil into each reactor. The soil was maintained at 70% to 80% of field capacity for the duration of the experiment. ^{14}C -pyrene was spiked into triplicate sets of reactors to determine the rate of mineralization to $^{14}\text{CO}_2$ as well as transformation of the parent compound as a function of oxygen concentration in the gas phase within the glass microcosms. The use of radiolabeled chemicals allowed a chemical mass balance to be generated that included fractions of radiolabeled carbon that were mineralized, volatilized, solvent extractable and soil bound. For each biometer flask, a carbon dioxide trap (10 ml of 0.1 N KOH) was used to collect $^{14}\text{CO}_2$ to determine the amount of mineralized ^{14}C -pyrene. A polyurethane plug, located in the side of each flask, was used for collection of volatile parent compounds and intermediates.

The desired O_2 concentrations were maintained in the reactors using premixed ratios of $\text{N}_2:\text{O}_2$ in the feed stock gases. Each reactor was flushed with the appropriate O_2 concentration three times per week. Flushing was accomplished using a vacuum system where 0.5 atmospheres were drawn out of the reactor and replaced with the premixed gas. This was repeated seven times. Measurements of the headspace oxygen concentrations in the reactors were done repeatedly using gas chromatography. Leaks were not found in the reactors, and oxygen concentrations were maintained at $\pm 1\%$ of the desired concentrations.

Experimental design

Triplicate reactors for each oxygen concentration were used for a total of 60 reactors, i.e., 3 replicates x 5 oxygen concentrations x 2 modes (non-poisoned and poisoned) x 2 times (day 0 and day 70) = 60 reactors. The reactors were incubated at the different oxygen concentrations for two to three weeks before the ^{14}C -pyrene was added in

order to ensure steady state with regard to oxygen concentration. Immediately before the radiolabeled pyrene was added, a set of reactors were extracted to obtain concentrations at the beginning of the experiment. Radiolabeled pyrene was added as well as the 10 ml of KOH, and reactors were monitored at seven day intervals for $^{14}\text{CO}_2$. The soil gas phase and KOH solution within the biometer flasks were accessible through two hypodermic needles with two-way Luer-Lock stop cocks attached. After 70 days, the production of $^{14}\text{CO}_2$ in the microcosms ceased to be statistically significant, and each reactor was destructively analyzed using the entire soil contents (50 grams).

For evaluation of the biological component of degradation, a triplicate set of reactors at each oxygen concentration was poisoned with mercuric chloride (HgCl_2) that served as controls to evaluate abiotic loss within the soil. The poisoning procedure consisted of mixing a solution of HgCl_2 and water with the soil to be poisoned and bringing it to a concentration of 1000 mg HgCl_2/kg dry weight of soil at a water content of 70% of field capacity.

Quality assurance/quality control

As part of the QA/QC plan, method blanks were extracted and analyzed as well as matrix spike duplicates following the guidelines given in Test Methods for Evaluating Solid Wastes, U.S. EPA SW-846 [26]. Method blank recoveries and matrix spike duplicate recoveries were in accordance with the data quality objectives outlined for this project. Method detection and quantitation limits were determined for the PAH using gas chromatography/mass spectrometry (GC/MS) analysis. The method quantitation limits for each PAH are as follows: naphthalene, 1.4 mg/kg; acenaphthylene, 1.1 mg/kg; acenaphthene, 1.3 mg/kg; fluorene, 1.5 mg/kg; phenanthrene, 2.1 mg/kg; anthracene, 1.7 mg/kg; fluoranthene, 2.8 mg/kg; pyrene, 2.8




Figure 2. Mineralization of ¹⁴C-pyrene in non-poisoned soil microcosms as a function of oxygen concentration. Values are the means for triplicate reactors.

mg/kg; benzo(a)-anthracene, 1.3 mg/kg; chrysene, 1.7 mg/kg; benzo(b)fluoranthene, 2.4 mg/kg; benzo(k)fluoranthene, 1.9 mg/kg; benzo(a)-pyrene 1.6 mg/kg; dibenzo(a,h)anthracene, 2.2 mg/kg; benzo(g,h,i)perylene, 1.6 mg/kg; and indeno(1,2,3-cd)pyrene, 1.7 mg/kg. Internal standards and surrogate standards for GC/MS were used and analyzed within the guidelines found in SW-846 [26]. All QA/QC for this experiment met the data quality objectives for the project.

Chemical analyses

The soil samples from the microcosms were extracted using soxhlet extraction, U.S. EPA Method 3540 [26]. Silica gel cleanup, U.S. EPA Method 3630 [26], was used to separate the PAH compounds from the heavy oils and greases. The instrumental analysis used to measure nonradiolabeled PAH in the soil was gas chromatography/mass spectrometry (GC/MS), U.S. EPA Method 8270 [26]. A portion of the extract was analyzed for ¹⁴C.

After the soil in the microcosms was extracted using the soxhlet procedure, the soil was allowed to air dry. After the soil was air dried, it was mixed thoroughly, and triplicate subsamples of soil from each sample were weighed out and combusted for determination of soil-bound ¹⁴C. The polyurethane plug was extracted using the soxhlet





Figure 3. Mineralization of ¹⁴C-pyrene in poisoned soil microcosms as a function of oxygen concentration. Values are the means for triplicate reactors.

extraction procedure, and the extract was also analyzed for ¹⁴C. All ¹⁴C analysis was performed on a Beckman LS 6000 Liquid Scintillation Counter.

RESULTS AND DISCUSSION


Results of the mineralization study for ¹⁴C-pyrene are plotted in Figures 2 and 3 for the non-poisoned and poisoned microcosms, respectively. Each data point plotted represents the mean of triplicate measurements. Results of ¹⁴CO₂ measurements for ¹⁴C-pyrene in poisoned reactors (Figure 3) show less than 0.5% mineralization at any soil gas oxygen concentration. This trend is consistent throughout the 70-day incubation period, indicating that biodegradation was responsible for the mineralization of ¹⁴C-pyrene.

Results shown in Figure 2 indicate that mineralization increased significantly as oxygen concentrations in the soil gas phase were increased above zero percent in the non-poisoned microcosms. Both rates and extents of mineralization were similar for all non-zero oxygen concentrations over the 70-day incubation. Ranges for cumulative mineralization of added ¹⁴C-pyrene after 70 days of incubation were 45% to 55% within the oxygen concentration range of 2% to 21%. Although mineralization was measured as 13% after 70 days incubation at 0% oxygen, the stan-



dard deviation was 12.86%. Mineralization in the microcosms incubated at 2%, 5%, 10% and 21% oxygen were not significantly different until day 56. At day 56, the mineralization in 2% oxygen samples became significantly larger than the mineralization in the 21% oxygen samples. Significant differences were determined using the least significant difference (LSD) obtained from the analysis of variance (ANOVA) method. Mineralization in the 5% and 10% microcosms was not significantly different from either the 2% or 21% samples throughout the duration of the experiment. At 0% oxygen concentration in the soil gas phase, mineralization of ^{14}C -pyrene was found to be statistically insignificant.

Figure 4 shows results of a two way analysis of variance for oxygen concentrations and percent mineralization for non-poisoned reactors averaged over the 70-day incubation period. Figure 4 indicates that average mineralization was similar at 35% to 40% for all non-zero oxygen concentrations in the soil gas phase and significantly less for reactors incubated at zero oxygen concentration. The least significant difference was 2.39. Using this LSD, average mineralization at the atmospheric oxy-



gen level is significantly lower than at the 2%, 5% and 10% levels. This analysis was not appropriate for poisoned reactors because there was no significant mineralization observed at any soil gas oxygen concentration.

Results of the mineralization evaluation indicate that even small soil gas oxygen concentrations substantially below atmospheric oxygen levels are sufficient to allow mineralization of pyrene in the contaminated soil evaluated. An explanation of why mineralization was less at 21% oxygen than at 2%, 5% and 10% oxygen levels is not available at this time. In water saturated marine environments and in uncontaminated saturated soil, it has been shown that pyrene and other PAHs will biodegrade at low oxygen concentrations [7, 27, 28].

The Michaelis-Menten relationship for enzyme kinetic analysis was used to determine biokinetic parameters including maximum mineralization rate (R_{\max}) and half-saturation constant (K_m). Figure 5 is a plot of the results showing mineralization rate as a function of oxygen concentration in the soil gas. From this plot, the R_{\max} value was found to be approximately 1.46% mineralization/day. The half saturation constant (K_m) was determined to be 1% oxygen concentration in the soil gas phase. Using Henry's law of partial pressures, the equilibrium concentration of oxy-

gen in the soil water phase was calculated to be 0.44 mg/L.

Results for the chemical mass balance on ¹⁴C-pyrene after 70 days of incubation are presented in Tables 2 and 3 for the non-poisoned and poisoned reactors, respectively. Results presented in these tables indicate that in every case the total mean recovery of ¹⁴C from the microcosms was above 85% for all mass balances conducted. With regard to mineralization, values at the termination of the incubation period are provided, and they indicate that mineralization was the most significant pathway for pyrene transformation, with the exception of microcosms incubated at 0% oxygen concentration, where mineralization was statistically equivalent to zero. Soil extractable ¹⁴C was the second most significant fraction with a range of 20 to 25% recovery for non-zero oxygen concentrations. For the soil incubated at 0% oxygen concentration, solvent extractable ¹⁴C accounted for the most significant fraction of

the ¹⁴C. Soil extractable ¹⁴C could include any ¹⁴C that partitions into the solvent phase and may include the parent compound as well as solvent soluble intermediates. The solvent extractable phase was not evaluated further.

The third most significant fraction of ¹⁴C in the non-poisoned reactors was the soil bound fraction—about 15% ¹⁴C recovery for non-zero oxygen reactors and about 8% for 0% oxygen reactors. Results indicate that in the presence of oxygen a larger fraction of the ¹⁴C was incorporated into the soil phase to the extent that it could not be extracted with the solvent. This may be due to chemical binding of intermediates to soil organic matter generated in the biodegradation of pyrene in a process referred to as humification [23].

Regarding volatilization of ¹⁴C, results in Table 2 indicate that less than 0.3% ¹⁴C was volatilized over the 70-day incubation period.

Oxygen Concentration	% Mineralized	% Volatilized	% Soil Extractable	% Soil Bound	% ¹⁴ C Mass Recovered
0%	12.96±12.86	0.27±0.03	69.68±14.03	8.47±0.8	91.39±19.05
2%	53.95±4.76	0.10±0.08	22.32±4.59	15.08±2.40	91.46±7.03
5%	51.99±1.21	0.06±0.02	20.74±1.21	15.66±1.11	88.45±2.04
10%	50.92±1.45	0.14±0.14	20.47±0.69	14.20±1.29	85.73±2.06
21%	45.51±1.02	0.08±0.01	24.66±1.54	15.41±0.77	85.66±2.00

Table 2. Average distribution of ¹⁴C in non-poisoned microcosms spiked with ¹⁴C-pyrene ± standard deviation.

Oxygen Concentration	% Mineralized	% Volatilized	% Soil Extractable	% Soil Bound	% ¹⁴ C Mass Recovered
0%	0.06±0.01	0.35±0.04	86.01±0.90	8.56±0.64	94.98±1.12
2%	0.09±0.01	0.22±0.05	82.01±0.59	8.77±0.52	91.09±0.79
5%	0.09±0.01	0.29±0.06	77.78±0.73	10.97±1.71	89.13±1.86
10%	0.09±0.01	0.29±0.08	76.79±1.46	12.46±1.40	89.63±2.02
21%	0.08±0.02	0.36±0.02	87.93±10.31	8.34±0.37	96.71±10.32

Table 3. Average distribution of ¹⁴C in poisoned microcosms spiked with ¹⁴C-pyrene ± standard deviation.




Figure 6. Nonradiolabeled pyrene concentrations in non-poisoned reactors at day 0 and day 70 as a function of soil gas oxygen concentration.




Figure 7. Benzo(a)anthracene concentrations in non-poisoned reactors at day 0 and day 70 as a function of soil gas oxygen concentration.

Results presented in Table 3 for poisoned microcosms indicate insignificant mineralization and volatilization, with the major fraction of ^{14}C measured as soil extractable ^{14}C within the range of 77% to 88%. This high value for solvent recovery is likely due to the absence of biotic transformations of the ^{14}C -pyrene in the soil. Soil bound ^{14}C was observed to be within the range of 8% to 12%. There were no differences between observations at non-zero oxygen concentrations and 0% oxygen concentrations. Total mean ^{14}C mass balances for the poisoned microcosms were all above 85% as observed with the non-poisoned microcosms.

Initial and final PAH concentrations were determined for all 16 priority PAH compounds in both the poisoned and non-poisoned microcosms. Eight of the 16 PAH compounds were found to be below the method quantitation limit (MQL) discussed in the Experimental Design section. Because the concentrations of these compounds were below the MQL, statistical analysis could not be conducted with good reliability; therefore, results for the degradation of these compounds are not reported. Eight PAH compounds were found in concentrations above the method quanti-

tation limit for both poisoned and non-poisoned reactors at day 0 and day 70. These eight compounds were: naphthalene, anthracene, pyrene, benzo(a)anthracene, chrysene, benzo(b)fluoranthene, benzo(k)fluoranthene and benzo(a)pyrene.

An analysis of variance (ANOVA) was used to determine whether different factors such as time and oxygen concentration had a significant effect on PAH loss in the non-poisoned reactors. Three 4-ring compounds (pyrene, benzo(a)anthracene and chrysene) were selected for the evaluation with regard to the effect of soil gas oxygen. Figures 6-8 are graphs of this response. For pyrene (Figure 6), there are significant differences at day 70 between the 0% oxygen level and the 5% and 21% oxygen levels for loss of pyrene. For benzo(a)anthracene (Figure 7) there is significant difference between 0% oxygen and the 2%, 5% and 21% oxygen levels. Results indicate a larger degree of biodegradation when oxygen is present compared to when it is not present.

There was also a statistically significant difference in the degree of degradation of




Figure 8. Chrysene concentrations in non-poisoned reactors at day 0 and day 70 as a function of soil gas oxygen concentration.

chrysene as a step-function of oxygen. Results shown in Figure 8 indicate that the addition of oxygen to the soil gas phase caused a statistically significant loss of chrysene over 70 days; however, when no oxygen was available the concentration of chrysene did not significantly change.

There were also significant differences between PAH concentrations in the poisoned and non-poisoned microcosms for all eight compounds at all non-zero oxygen concentrations. These results are given in Table 4. From this table, it is shown that

when the soil was not poisoned there was a significant loss of PAH compared to the poisoned soil when oxygen was present. The results for the nonradiolabeled PAH agree with ¹⁴C-pyrene results with regard to greater reduction in PAH concentration measured in non-poisoned non-zero oxygen microcosms.

CONCLUSIONS

Results using ¹⁴C-pyrene as an indicator of the amount of degradation that occurs under different oxygen concentrations demonstrated that oxygen tension does have a significant effect on rate and amount of degradation of ¹⁴C-pyrene. At oxygen concentrations of 2% to 21%, mineralization accounted for between 45% and 55% of the ¹⁴C-pyrene spiked onto the soil after 70 days. At these non-zero soil gas oxygen concentrations, mineralization of ¹⁴C-pyrene was statistically significantly higher than at the 0% oxygen concentration, while mineralization of ¹⁴C-pyrene was not significant at the 0% oxygen level. Based on the negligible amount of mineralization measured in poisoned microcosms, mineralization that occurred in the non-poisoned microcosms was due to biological processes. For mineralization of ¹⁴C-pyrene, the half saturation constant (K_m) was calculated to be 1% oxygen concen-

PAH Compounds	Poisoned	Non-Poisoned	% Loss	LSD
Naphthalene	7.3	6.85	6.2	0.37
Anthracene	23.13	11.96	48.3	3.28
Pyrene	15.73	6.72	57.3	1.73
Benzo(a)anthracene	4.88	2.66	45.5	0.45
Chrysene	8.06	4.73	41.3	0.76
Benzo(b)fluoranthene	8.9	7.31	17.9	0.92
Benzo(k)fluoranthene	8.39	6.74	19.7	0.88
Benzo(a)pyrene	11.06	9.31	15.8	1.21

¹Eight PAH compounds were found below the MQL and therefore were not quantified.

Table 4. Poisoned and non-poisoned microcosms PAH concentrations (mg/kg), percent difference and least significant difference averaged over time and non-zero oxygen levels.¹

tration, and the maximum mineralization rate (R_{\max}) was 1.46% mineralization/day.

Based on a chemical mass balance and analysis for ^{14}C , biological mineralization was the dominant fate mechanism at all non-zero oxygen concentrations. Soil extractable ^{14}C was the second largest fraction of recovered ^{14}C from microcosms incubated at non-zero oxygen concentrations, and was the dominant fate of ^{14}C -pyrene incubated at 0% oxygen concentrations in non-poisoned microcosms and in all poisoned microcosms. Soil bound (non-solvent extractable) ^{14}C was the third largest fate mechanism at all non-zero oxygen concentrations. Volatilization was minimal for all oxygen concentrations evaluated.

A recommendation for soil gas oxygen would be a minimum of 2% to 5% in buried lifts of a prepared bed bioreactor in order to maintain conditions conducive for continued biodegradation of pyrene and other PAH. Low oxygen concentration requirements indicate minimal pumping of oxygen through the buried lifts would be required and then only when the soil gas oxygen concentration decreases to below 2% to 5%. If these critical oxygen levels necessary for biodegradation of PAH are present, then continued biodegradation of target chemicals in the lower lifts may allow the lower lift to reach target remediation levels after an upper lift has been applied. Treatment time may be shortened by placing a new contaminated soil lift on top of a lower lift before the soil comprising the lower lift reaches target remediation levels. Results from this research will be used to determine the potential for decreasing the required time for treatment of contaminated soil in prepared bed systems by maintaining the necessary concentration of oxygen in the soil gas phase for active aerobic biodegradation of PAH in buried lifts within the prepared bed.

ACKNOWLEDGMENTS

The authors wish to acknowledge the significant support of the Champion International staff at Libby, Montana. Without the support of these individuals, this project could not have been accomplished. Also, we would like to express thanks to Neal Yancey and Hong Shang at the Utah Water Research Laboratory, Logan, Utah, for their technical assistance. This research was funded by the Great Plain/Rocky Mountain Hazardous Substance Research Center in conjunction with the U.S. EPA, through the Robert S. Kerr Environmental Research Laboratory, (RSKERL) Ada, OK. Support was provided to Utah State University, Logan, UT. Dr. Ronald C. Sims is the principal investigator and Scott Huling is the U.S. EPA project officer.

REFERENCES

1. S.E. Herbes, G.R. Southworth and C.W. Gehra, Organic Contamination in Aqueous Coal Conversion Effluents: Environmental Consequences and Research Priorities, In: D.D. Hemphill (Eds.), Trace Substances in the Environment: A Symposium, Univ. of Missouri, Columbia, MO, 1976.
2. R.J. Mahmood, Enhanced Mobility of Polynuclear Aromatic Hydrocarbons in Unsaturated Soils, Utah State University, Ph.D. Dissertation, 1989.
3. R.C. Sims and M.R. Overcash, Fate of Polynuclear Aromatic Compounds (PNAs) in Coal Systems, Residue Reviews, 88 (1983) 1-68.
4. G.D. McGinnis et al., Characterization and Laboratory Soil Treatability Studies for Creosote and Pentachlorophenol Sludge and Contaminated Soil, U.S. EPA, Washington D.C., EPA 600/S2-88/055, 1988.

5. R.M. Atlas, Microbial Degradation of Petroleum Hydrocarbons: An Environmental Perspective, *Microbiological Reviews*, 45 (1981) 180-209.
6. H. Kobayashi and B.E. Rittman, Microbial Removal of Hazardous Organic Compounds, *Env. Sci. & Tech.*, 16 (1982) 170A-183A.
7. H.S. Park and R.C. Sims, Biodegradation of Polycyclic Aromatic Hydrocarbon Depending Upon Oxygen Tension in Unsaturated Soil, In: P.N. Cheremisinoff and Y.C. Wu (Eds.), *Hazardous Waste Management Handbook: Technology, Perception, and Reality*, Prentice-Hall Inc., Englewood Cliffs, NJ, 1994, pp. 257-273.
8. J.W. Foster, Bacterial Oxidation of Hydrocarbons, In: O. Hayaishi (Ed.), *Oxygenases*, Academic Press, New York City, NY, 1962.
9. C.E. Zobell and J.F. Prokop, Microbial Oxidation of Mineral Oils in Barataria Bay Bottom Deposits, *Z. Allg. Mikrobiol.*, 6 (1966) 143-162.
10. I.J. Higgins and J. Quayle, Oxygenation of Methane by Methane Grown *Pseudomonas methanica* and *Methanomonas methanooxidas*, *J. Biochem.*, 118 (1970) 201-208.
11. S. Dagley, Microbial Degradation of Organic Compounds in the Biosphere, *American Scientist*, 63 (1975) 681.
12. R.W. Hansen and R.E. Kallio, Inability of Nitrate to Serve as a Terminal Oxidant for Hydrocarbons, *Science*, 125 (1957) 1198-1199.
13. J.W. Foster, Hydrocarbons as Substrates for Microorganisms, *Antonie van Leeuwenhoek, J. Microbiol. Serol.*, 28 (1962) 241-27.
14. J.J. Perry, Microbial Cooxidation Involving Hydrocarbons, *Microbiological Reviews*, 43 (1979) 59-72.
15. C.S. Criddle, L.A. Alvarez and P.L. McCarty, Microbial Processes in Porous Media, In: J. Bear and M. Yavuz Corapcioglu (Eds.), *Transport Processes in Porous Media*, Kluwer Academic Publishers, Netherlands, 1991, pp. 639-691.
16. J.L. Sims, R.C. Sims and J.E. Matthews, Bioremediation of Contaminated Surface Soils, U.S. EPA, Washington D.C., EPA 600/9-89/073, 1989.
17. D.F. Pope and J.E. Matthews, Bioremediation Using the Land Treatment Concept, U.S. EPA, Washington D.C., EPA 600/R-93/164, 1993.
18. S.G. Huling, D.F. Pope, J.E. Matthews, J.L. Sims, R.C. Sims and D.L. Sorensen, Land Treatment and the Toxicity Response of Soil Contaminated with Wood Preserving Waste, *Remediation*, Spring, 1995.
19. H.L. Penman, Gas and Vapor Movements in the Soil, *J. Agr. Sci.*, 30 (1940) 570-581.
20. L.H. Stolzey and J. Letey, Characterizing Soil Oxygen Conditions with Platinum Electrode, *Advances in Agronomy*, 16 (1964) 249-279.
21. Y.S. Lin, Biodegradation of Organic Hazardous Wastes by Land Treatment Process, thesis presented to the Oklahoma State University, Norman, Oklahoma, Ph.D. Dissertation, 1986.
22. R.C. Sims, D.L. Sorensen, J.L. Sims, J.E. McLean, R. Mahmood, R.R. Dupont, J. Jurinak and K. Wagner, *Contaminated Surface Soils In-Place Treatment Techniques*, Noyes Publications, Park Ridge, New Jersey, 1996.

23. R.C. Sims, J.L. Sims, D.L. Sorensen and J.E. McLean, Champion International Superfund Site, Libby, Montana: Bioremediation Field Performance Evaluation of the Prepared Bed Land Treatment System, 1995, U.S. EPA (in press; Mr. Scott Huling, Robert S. Kerr Environmental Research Laboratory, Ada, OK, U.S. EPA Project Officer).
24. M.D. Lee, Biodegradation of Organic Contaminants in the Subsurface of Hazardous Waste Sites, Rice University, Ph.D. Dissertation, 1986.
25. L.D.A. Stetzenbach, The Degradation and Utilization of Polycyclic Aromatic Hydrocarbons by Indigenous Soil Bacteria, The University of Arizona, Ph.D. Dissertation, 1986.
26. U.S. EPA, Test Methods for Evaluating Solid Waste, Physical/Chemical Methods, 3rd ed., SW-846, U.S. Environmental Protection Agency, Washington, D.C., 1986.
27. J.E. Bauer and D.G. Capone, Degradation and Mineralization of Polycyclic Aromatic Hydrocarbons Anthracene and Naphthalene in Intertidal Marine Sediments, *Applied and Environmental Microbiology*, 50 (1985) 81-90.
28. C. Fritzsche, Degradation of Pyrene at Low Defined Oxygen Concentrations by a *Mycobacterium* sp., *Applied and Environmental Microbiology*, 60 (1994) 1687-1689.