

Greenhouse Gas Accounting in Agriculture:

Dairy Industry Experience

Greg Thoma

Darin Nutter

Rick Ulrich

Dae Soo Kim

College of Engineering, University of Arkansas

Marty Matlock

Jennie Popp

Nathan Kemper

Zara Niederman

UA Center for Agricultural and Rural Sustainability

David Shonnard

Felix Adom


Charles Workman

Chemical Engineering, Michigan Technological University



Why an LCA of Milk Production?

- Consumers are interested in product sustainability.
 - Increasing willingness to make purchase decisions based on (perceived) environmental impacts.
 - Potential for reduction in consumption of dairy.
- Science-based LCA can provide:
 - Data that will allow the industry to identify and engage more sustainable approaches
 - Reduce environmental impacts that can be validated through LCA measurements.
- Other benefits:
 - Systems understanding allows positioning dairy products in the marketplace based on sustainable attributes respond proactively to consumer concerns.
 - Supports industry's ability to work with retailers to educate consumers about agricultural and food sustainability issues.
 - Establish a baseline for GHG offset projects which may result from legislation in the future



LCA Methodology

ISO 14044 compliant

Goal: Determine GHG emissions associated with consumption of one gallon of milk to US consumer.

Scope: Cradle to grave. Specifically including pre-combustion burdens for primary fuels and disposal of packaging.

National Scale Analysis – drives data collection



Global Warming Potential of Fluid Milk

Scope: grass to glass



Life Cycle Inventory – Data Drives the Work

Surveys:

- 1) Dairy Producer (~535; 9% response rate)
- 2) Farm to processor transportation data
(~150,000 round trips – 2007 only)
- 3) Milk Processor (50 plants responded)

Published Literature:

- 1) Peer Reviewed Literature
 - a) Enteric Methane, Nitrogen and Methane from manure management
 - b) Life cycle inventory data for crop production (NASS, Budgets, USLCI)
- 2) Other Publications (e.g. IPCC, EPA)
- 3) Expert opinion




Dairy Feed: crop production

Fertilizer, pesticides, fuel & processing



What Cows Eat



Calculating Feed Footprint:

Data Sources

- NASS data on fertilizers, fuels, pesticides and other inputs
 - States with complete data averaged
 - CO₂e burden of inputs accounted
 - Production burdens and N₂O from field accounted
- Production budgets from state Ag extension agents used for silage, hay, and pasture
 - Budget prices converted to estimated quantities
 - Similar accounting: state budgets aggregated and CO₂e accounted



Data Challenges

- Incomplete data sets
 - Fertilizer type by crop, pesticides, fuel use
- Complexity of dairy rations
 - Lack of extant LCA for many items
 - Almond hulls, citrus pulp, apple pomace
 - Surrogate feeds adopted
- Crop production practices
 - Tillage, soil type, etc affect N_2O
 - Lime fate (soil acidity)
- Consistency of background data
 - System boundaries, linking to relevant upstream information




Challenges: Allocation

- System expansion
- Physical causality
- Economic value
- Mass/Energy content
- By-products:
 - Distiller's grains, grain meals, pulp, etc
- Milk and beef
- Cream and milk products
- Refrigeration at retail and in-home



The Feedprint



National Feed Carbon Footprint, Fractional Contribution by Feed Type

Dairy Fluid Milk

Farm Level Data Collection

Producer Survey for On-Farm Data






Collect representative data from U.S. dairy producers to establish a carbon footprint baseline relevant to conditions in the United States.

Single farm footprint will have similar goal, but obviously require site specific data and understanding of practices

Producer Survey

- 43 questions in 9 areas
 - About Your Facility
 - On-Facility Crop Production
 - Manure Management
 - Energy Usage
 - Housing & Milking Information
 - Animal Feedstuffs & Grazing Practices
- 500+ usable surveys returned



Producer Survey Data

- Tremendous effort from farmers
- Generally very conscientious in providing information
 - Despite multiple revisions, still some confusion
 - Missing information
 - Herd demographics often incomplete
 - Manure management quantities inconsistent
 - On-farm crop production data
 - Animal diets – information runs the gamut!
- Lesson:
 - Farm scale analysis will need knowledgeable data collectors and very patient farmers



Data Management & Reconciliation

- Each survey entered twice; copies compared and corrected by reference to original
- Outliers identified
 - (eg 13 day calving interval → months)
 - Milk production
- Missing data interpolated by substitution with weighted regional average
 - Herd demographics
 - Dry matter intake (regional average ration used when missing)
 - MUN, fat and protein



On-Farm Emissions

Enteric, manure, energy


Data from producer survey & existing models



Diet and Enteric Methane

- Diet from survey matched with NRC feed database to determine relevant parameters:
 - Crude Protein, DE, NE_g , NE_L , ...
 - Milk to Beef allocation calculations
- Model Comparisons:
 - Ellis DMI model has lowest RMSE
- Biogenic GHGs
 - Methane explicitly accounted
 - Other biogenic carbon cycling is not explicitly accounted
 - In line with current IDF discussions






Milk / Beef Allocation Choices

- System expansion
 - Beef from dairy not quite equivalent to 'Angus'
 - Culled cows similar to breeder stock; bull calves have no real equivalent
 - LCA on US beef has not been completed
- Biological / Causal
 - Each feed has different conversion efficiency
 - Determine digestible energy consumed for:
 - Growth
 - Milk production
 - Determine total feed energy for milk and beef, use this to define allocation fraction – Then all emissions allocated with this ratio
- Economic Value




Schematic of energy flow accounting for allocation




Cradle to Farm Gate Results




National Farm Gate GWP for US Milk



(Allocation OFF) Sensitivity Analysis: Slurry with Crust, 2 Months on Pasture



Sensitivity Analysis: Slurry with Crust, 2 Months on Pasture



Farm Gate through Consumption

Highlights




Transportation of Raw Milk from Farms to Processing Plants

- ~150,000 round trips (6000 gallon trucks) in 2007
 - Combined data from several dairy cooperatives.
 - – 11% of fluid milk delivered in 2007
-
- Contained in this data for each trip:
 - Day of the year that the trip started, from 1 to 365, in 2007
 - Latitude and longitude of the plant
 - Latitude and longitude of each farm
 - Pounds of milk picked up at each farm



Cumulative Histogram of Trip Lengths

average =
490 miles





Milk Processing & Distribution

Processing (pasteurization),
packaging, and distribution to retail




Processor Survey Details

- Total of 50 processing plants provided data for 2007
 - ~ 25% of all fluid milk processed in 2007
- Data verification and analysis
 - Two clarification rounds
 - Detailed blow molding equipment specifications



Processing Contribution to GWP



Retail allocation

$$\frac{\left(\frac{\text{kW-hr}}{\text{month}} \text{ to refrigerators_containing_milk} \right) \left(\frac{\text{kg CO}_{2e}}{\text{kW-hr}} \right) \left(\frac{\text{mass milk in refrigerators}}{\text{mass total in refrigerators}} \right)}{\left(\text{fraction of milk used by buyer} \right) \left(\frac{\text{kg}}{\text{month}} \text{ milk sold} \right)}$$

+

$$\frac{\left[\left(\frac{\text{kW-hr}}{\text{month}} \text{ to store} \right) - \left(\frac{\text{kW-hr}}{\text{month}} \text{ to all refrigerators} \right) \right] \left(\frac{\text{kg CO}_{2e}}{\text{kW-hr}} \right) \left(\frac{\text{milk sales}}{\text{total sales}} \right)}{\left(\text{fraction of milk used by buyer} \right) \left(\frac{\text{kg}}{\text{month}} \text{ milk sold} \right)}$$


+

$$\frac{\left(\frac{\text{kg refrigerant}}{\text{month}} \right) \left(\frac{\text{kg CO}_{2e}}{\text{kg refrigerant}} \right) \left(\frac{\text{mass milk in refrigerators}}{\text{mass total in refrigerators}} \right)}{\left(\text{fraction of milk used by buyer} \right) \left(\frac{\text{kg}}{\text{month}} \text{ milk sold} \right)}$$


$$= \text{kg CO}_{2e} / \text{kg milk consumed}$$



Retail/Consumer



End-of-Life (Packaging disposal)




LCA Software Tools

- Excel & VBA
- MatLab
- SimaPro 7.1 (Pre Consultants)
 - Databases of LCI data
 - Eco Invent
 - US LCI
 - Franklin Associates
- EIO LCA <http://www.eiolca.net>
- Open IO – Carbon <http://www.open-io.org>
- Earthster <http://www.earthster.org>
 - Open source data repository and computational engine




Supply Network



Why an LCA of Milk Production?

- Consumers are interested in product sustainability.
 - Increasing willingness to make purchase decisions based on (perceived) environmental impacts.
 - Potential for reduction in consumption of dairy.
- Science-based LCA can provide:
 - Data that will allow the industry to identify and engage more sustainable approaches
 - Reduce environmental impacts that can be validated through LCA measurements.
- Other benefits:
 - Systems understanding allows positioning dairy products in the marketplace based on sustainable attributes respond proactively to consumer concerns.
 - Supports industry's ability to work with retailers to educate consumers about agricultural and food sustainability issues.
 - Establish a baseline for GHG offset projects which may result from legislation in the future



Whose carbon is it?

- ISO compliant LCA should be cradle to grave in scope
 - Does the dairy farmer get credit for more efficient corn production?
 - **Yes:** in the context of a full LCA accounting of the entire supply chain, they can legitimately claim milk has overall lower impact.
 - *AND*
 - **No:** in the context of carbon credits or carbon trading. Here the WRI concepts of Scope 1,2,3 are useful.
 - Scope 1: on-site emissions (enteric methane, diesel combustion)
 - Scope 2: indirect emissions (mostly electricity)
 - Scope 3: indirect emissions further up the supply chain (N₂O from fertilizer manufacture, or diesel combustion for commodity crop production)
- LCA is a useful tool and framework for Scope 1 assessment, but for carbon trading full LCA is not really appropriate



Summary

- Bottom line: Milk LCA results
 - Add up all the emissions → 1 gallon of milk consumed in the US is less than equivalent to burning 1 gallon of gasoline → driving 20 miles.
- GHG Accounting in Agriculture
 - Complex systems – crops, soil, climate interactions
 - Data quality is likely to remain a concern – e.g., how much methane is really released at a particular farm?
 - Allocation can significantly affect results
 - Sensitivity analysis can highlight areas where good measurements are critical to proper assessment of the CF



Acknowledgements

Dairy Management, Incorporated
and
The Innovation Center for US Dairy

