
CALCIUM AND MAGNESIUM INTERFERENCE STUDIES FOR THE BINDING OF HEAVY METAL IONS IN SOLUTION BY *MEDICAGO SATIVA* (ALFALFA)

J.L. Gardea-Torresdey^{*1}, K.J. Tiemann¹, J.H. Gonzalez¹, J.A. Henning², and M.S. Townsend²,
¹Department of Chemistry, The University of Texas at El Paso, El Paso, TX, 79968, Phone: 915-747-5359, FAX: 915-747-5748, and ²Department of Agronomy & Horticulture, New Mexico State University, Las Cruces, NM, 88003

ABSTRACT Previous batch laboratory experiments performed to determine the potential ability of seven different varieties of *Medicago sativa* (alfalfa) revealed that the African shoots population was able to efficiently bind copper(II) and nickel(II) from aqueous solutions. Batch laboratory interference studies were performed with various calcium and magnesium concentrations (0.1 mM to 1 M) in order to ascertain the effects of these ions on the heavy metal binding ability of African alfalfa shoots. Results from these studies have shown that calcium and magnesium did not seriously reduce the binding of copper(II) and lead(II) to African alfalfa shoots. However, high concentrations of calcium and magnesium significantly reduced chromium(III), cadmium(II), nickel(II), and zinc(II) binding to African shoots. In addition, all these experiments were repeated maintaining the ionic strength constant, and similar results were obtained. Interference studies were also conducted in order to determine the effects of hard cations under flow conditions with silica-immobilized African alfalfa shoots. The information obtained from these studies will be useful for an innovative method of heavy metal ion removal and recovery from contaminated waters.

KEYWORDS: bioremediation, phytofiltration, alfalfa, *Medicago sativa*, interference, heavy metal binding

INTRODUCTION

Due to the increase in industrial activity, an alarming amount of toxic heavy metals have been released into the environment endangering natural ecosystems and public human health [1-3]. Industries such as smelters, metal refineries, and mining operations have been indicated as major sources of metal release into the environment [4-8]. The United States has invested billions of dollars attempting to clean heavy metal-polluted ground waters and soils. However, recent studies have shown that many of the treated ground water sites have not been restored back to drinking water standards [9]. Current technologies that are employed are not only costly and

inefficient, but they also increase the contaminant exposure to cleanup crews. The limitations of conventional ground water remediation have spurred many investigations into alternative methods which are more cost effective. One of these technologies is bioremediation, the use of biological systems such as those employed to remediate petroleum. Bioremediation can change the offending organic contaminant into substrates such as CO₂ [10]. The problem existing with bioremediation and bioaccumulation of toxic metals is that although the metals' valence or oxidation state may be biologically converted, the metal is still present and poses an environmental threat. Many bioremediation

*To whom correspondence should be addressed (jgardea@utep.edu).

techniques have been studied as safer, more conventional technologies, yet these processes are not universally understood nor accepted. This is in part due to the fact that these processes rely on microorganisms which precipitate or solubilize the metal ions [11-13].

An innovative technology that is gaining momentum in the environmental field is phytoremediation. Phytoremediation is the use of plants to remove toxic contaminants from the environment. J.L. Schnoor, *et al.*, have proven in several applications that hybrid poplar trees have the ability to phytoremediate contaminated soils at a low cost [14]. W.F. Mueller and co-workers also have shown the potential use of *Datura innoxia* and *Lycopersicon peruvianum* to biotransform TNT wastes [15]. Many studies such as those conducted by Ernst, *et al.*, have demonstrated the heavy metal resistance in plants [16]. Only recently has the value of heavy metal accumulating plants been realized in the process termed phytoextraction to remove the metals from the soils. Raskin, *et al.*, have proven that crops such as *Brassica junica* (L.) can possess great potential for phytoextraction of various metals from soils [17, 18]. In addition, many other investigators have shown metal uptake by cultivars such as lettuce, corn, and mushrooms [19-22]. Alfalfa (*Medicago sativa*) has also shown to be extremely resistant to high levels of contaminants as well as a bioaccumulator [23-25]. Only within the last few years has this technology been implemented for cleaning up heavy metal contaminated waters. I. Raskin and many other investigators have proven that rhizofiltration is a practical way to remove toxic heavy metals from ground waters [26-29]. Although these are cost-effective alternatives to the more traditional methods, rhizofiltration takes time and poses a threat

to the cleanup crews who must transport the metal-accumulated plants.

Since plants have shown to be accumulators of heavy metals, a more practical application would be “phytofiltration”: the use of plant material to accumulate toxins in a filtration apparatus. Previous studies such as those conducted by Ramelow, Rayson, and Volesky as well as other investigators indicate that concentrating heavy metals with an immobilized biomass in a column form would be practical and cost effective [30-33]. Gardea-Torresdey, *et al.*, have shown alfalfa as a potential source for “phytofiltration” [34-37]. Several batch laboratory experiments determined that alfalfa possess the ability to bind various heavy metal ions from aqueous solutions. In addition, considerable amounts of the bound metal ions were recovered from the reusable immobilized biomaterial. This biorecovery process for toxic metals may be employed for a cheaper and more efficient strategy to recycle waste metals. At least one problem that arises with most metal filtration systems is the presence of hard cations such as magnesium and calcium in contaminated waters. These cations have been reported as high as 4,000 ppm in some of the Superfund sites in Texas. These hard cations usually clog conventional ion-exchange-type filtration systems.

The objective of our study is to determine the effects of various concentrations of calcium and magnesium on the ability of immobilized alfalfa to remove different metal ions from aqueous solutions. Several batch laboratory experiments were carried out with various concentrations of calcium and magnesium while keeping the studied metal ion concentration constant. Column experiments were performed under similar conditions to determine the effects of

calcium and magnesium under flow conditions.

METHODOLOGY

Alfalfa collection

Previous studies were performed with seven alfalfa populations which were selected as representatives from the many different varieties of alfalfa due to their individual characteristics. The populations studied include the alfalfa basic germplasms (African, Peruvian, Flemish, Ladak) and two cultivars (Malone, Moapa 69) which were obtained from plots that had received irrigation every two weeks during the growing season. In addition, one cultivar (Cal West 630) was examined from a dryland test, which received no irrigation. These studies have shown that some populations bind metal ions better than others, and the best selection was used for further studies. The African germplasm bound both copper and nickel very well and was chosen as the biomass source for this study.

The alfalfa tissues were collected from field studies conducted by Dr. John Henning and Shawn Townsend at New Mexico State University near Las Cruces, New Mexico. The plants were removed from the soil, washed, and the roots were separated from the shoot material (stems and leaves). All samples were oven dried at 90°C, then ground to pass through a 100-mesh screen using a Wiley mill.

Batch laboratory interference experiments for metal binding

These studies were conducted to determine the interference of calcium and magnesium ions with the binding of metal ions to African shoots. For each experiment, 300 mg of African shoot biomass were washed

twice with 0.01 M HCl by vortexing to remove any debris or soluble biomolecules that might interact with the metal ions. This was then followed by centrifugation to obtain a pellet after each wash. The washings were collected, dried, and weighed to account for any biomass weight loss. The washed biomass was resuspended in 60 ml of deionized water with tissue concentration approximately 5 mg per ml solution, and the pH of the solution was adjusted to 5.0. Two ml of the African shoot suspension (10 mg in 2 ml) was transferred into each of three test tubes. Solutions were prepared to include 0.1 mM metal ion solution at pH 5.0 for the following calcium and magnesium concentrations: 0.0 M, 0.1 mM, 0.2 mM, 1 mM, 2 mM, 10 mM, 20 mM, 0.1 M, 0.2 M, and 1 M. Because these hard cations coexist in the environment, these studies were also carried out with both the calcium and magnesium concentrations combined as well as separate to determine their effects. From each solution, 2 ml of the various concentrations of hard cations and 0.1 mM metal solutions were added to the three respective tubes and reacted with biomass. In addition, three controls containing the 0.1 mM metal of interest were retained at each of the various concentrations of calcium and magnesium. All the tubes were equilibrated on a rocker for one hour. The samples were then centrifuged at 3,000 rpm for five minutes, and the supernatants for the pellets were transferred to clean respective tubes. Final pHs for all tubes were recorded, and all metal analyses were performed by flame atomic absorption.

Immobilization of alfalfa biomass

In order to prevent clumping of the alfalfa cell material, it was necessary to use a support material composed of a polysilicate matrix. The ground African alfalfa shoots were immobilized following the method

described by Gardea-Torresdey, *et al.* [35]. The polymerized gel which fixed the plant material was oven dried overnight at 60°C and then ground and sieved to pass 20-40 mesh size.

Column experiments

One bed volume which consisted of 6 ml of immobilized biomass was packed into columns. The columns were conditioned by passing 10 bed volumes of 0.01 M sodium acetate buffer at pH 5.0. For each experiment, 120 bed volumes of 0.1 mM metal solution in 0.01 M sodium acetate at pH 5.0 was passed at a flow rate of 2 ml per minute for each of the various concentrations of calcium and magnesium. This experiment was repeated for 0.0 M, 0.1 mM, 10 mM, and 0.1 M solution concentrations of calcium and magnesium. Each bed volume was collected and analyzed by flame atomic absorption.

Recovery of adsorbed metal ions from immobilized alfalfa

To remove the bound metal ions being studied, 10 bed volumes of 0.1 M HCl were passed through the columns of immobilized African alfalfa shoots at a flow rate of 2 ml per minute. Each bed volume passed was collected and analyzed for metal content by flame atomic absorption. Calculations were performed to determine the percent of metal recovery from the amount of metal bound to the columns for each experiment.

Analytical procedure

Analyses for the metal ions studied were performed using a Perkin Elmer model 3110 Atomic Absorption Spectrometer with deuterium background subtraction. The methods and conditions followed for each metal analysis were obtained from the Perkin Elmer model 3110 Atomic

Absorption Spectrometer manual. Analytical wavelengths used for the various metals were as follows: cadmium—228.8 nm; chromium—358.2 nm; copper—327.4 nm; nickel—352.5 nm; lead—283.3 nm; and zinc—213.9 nm. Calibration of the instrument was performed within the range of analysis, and a correlation coefficient for the calibration curve of 0.98 or greater was obtained. Periodically the instrument response was checked throughout the analysis with known standards. Samples were read three times, and a mean value and relative standard deviation were computed. The difference between the initial control metal concentration and that observed in the effluent was assumed to be bound by the alfalfa biomass.

Electron microscopy

An Environmental Scanning Electron Microscope (ESEM) was used to gain insight into the biomass interaction with the polysilicate support matrix that was used. An Electroscan ESEM model 2020 was used to take micrographs of uncoated, oven-dried, silica-immobilized African alfalfa shoots in a water vapor atmosphere at three torr.

RESULTS AND DISCUSSION

Previous screening experiments performed to determine the copper and nickel binding characteristics of seven populations of *Medicago sativa* alfalfa indicated that the African germplasm was among the best candidates for further metal binding studies. African alfalfa shoots have been shown to bind cadmium(II), chromium(III), copper(II), nickel(II), lead(II), and zinc(II) in considerable levels. However, due to the presence of hard cations such as calcium and magnesium found in contaminated ground waters, batch laboratory interference studies were performed with African alfalfa shoots to determine the effects of the hard cations




FIGURE 1. EFFECTS OF CALCIUM ON METAL BINDING BY AFRICAN ALFALFA SHOOTS (CADMIUM(II) ■, CHROMIUM(III) ▲, COPPER(II), NICKEL(II) - - - -, LEAD(II) ●, ZINC(II) ◆).

on the binding of the above-mentioned metal ions.

Figure 1 is the data collected from the calcium interference study. As indicated on Figure 1, the amounts of lead(II) and copper(II) bound to the alfalfa biomass remain moderately constant, until concentrations of calcium exceed 20 mM. A small reduction in the binding was observed thereafter, but appreciable levels of lead(II) and copper(II) were still removed from solution with calcium concentrations as high as 1 M. These concentrations of calcium, which are nearly 40,000 ppm, are ten times the levels reported at some of the contaminated Superfund sites. The binding of cadmium(II), nickel(II), and zinc(II) were affected by increased concentrations of calcium more drastically than lead(II) and copper(II). Overall, there was nearly a 30% reduction in the binding ability of the African alfalfa shoots when exposed to the highest concentration of calcium. The ability to bind heavy metal ions at concentrations of 0.1 mM while being immersed in 10,000 times more concentration of calcium indicates that the African alfalfa shoots may




FIGURE 2. EFFECTS OF MAGNESIUM ON METAL BINDING BY AFRICAN ALFALFA (CADMIUM(II) ■, CHROMIUM(III) ▲, COPPER(II), NICKEL(II) - - - -, LEAD(II) ●, ZINC(II) ◆).

be selectively binding the heavy metals in solution. Therefore, specific binding sites with chemical functional groups that have higher affinities for heavy metals may be responsible.

Figure 2 represents the data observed from the magnesium interference studies. From Figure 2, it can be seen that there is nearly no effect of magnesium interference on the African alfalfa shoot's ability to bind copper(II) and lead(II). However, for cadmium(II), chromium(III), nickel(II), and zinc(II), a significant decline was observed. Overall, again there was nearly a 30% reduction in the binding ability of the African alfalfa shoots when exposed to the highest concentrations of magnesium.

Figure 3 illustrates the combined effects of calcium and magnesium upon African alfalfa shoots binding ability. Only a slight decline was encountered for copper(II) binding, and a reduction in lead(II) binding was observed again after 20 mM of calcium and magnesium were added. This reduction of




FIGURE 3. EFFECTS OF CALCIUM AND MAGNESIUM ON METAL BINDING BY AFRICAN ALFALFA SHOOTS (CADMIUM(II) ■, CHROMIUM(III) ▲, COPPER(II) ·····, NICKEL(II) - - - -, LEAD(II) ●, ZINC(II) ◆).

lead(II) and copper(II) binding follows the same trend as that seen in Figure 1. Since magnesium concentrations had little effect on copper(II) and lead(II) binding to the alfalfa biomass, this decrease should be due to the calcium concentrations alone. As seen in Figures 1 and 2, cadmium(II), chromium(II), nickel(II), and zinc(II) binding was reduced by the increasing concentrations of calcium and magnesium. An overall reduction of 40% of the heavy metal binding to the alfalfa biomass was observed. The combined effects of the calcium and magnesium at the highest levels only increased the overall interference by 10% rather than 30% as one would expect. This overall difference may indicate that it is the quantity of hard cations in solution rather than the identity of the hard ions that plays a role in the reduction of heavy metal binding to alfalfa biomass.

Previous batch laboratory experiments indicated that metal ion binding to biomass functional groups may occur via an ion-exchange type mechanism. Because the alfalfa biomass demonstrates selective binding, the interference observed may be due to the increased concentration of ions in solution, instead of competition for binding

sites. These batch interference experiments were carried out again, keeping the ionic strength constant, and similar results were seen as those found in Figures 1-3.

The occurrences of binding specific metal ions in an ion-rich solution might be explained by the binding constants of various ligands that could be responsible for the heavy metal binding. The binding constants for various functional groups such as carboxylates, sulfur groups, and amino groups have an overall higher binding affinity for the various metal ions studied rather than calcium and magnesium [38]. Because the binding constants for the heavy metal binding to the various functional groups have higher stability constants, it stands to reason that the metals would bind before the hard cations would. This would explain the specificity of the alfalfa binding.

In addition to performing batch laboratory interference studies, column experiments were conducted to determine whether various concentrations of calcium and magnesium would affect the heavy metal binding ability of African alfalfa shoots under flow conditions. In order to maintain optimal flow rates, a polysilicate support material was used to immobilize the alfalfa biomass. Figure 4 shows an environmental scanning electron micrograph obtained of the silica immobilized African alfalfa shoots. As observed in Figure 4, a piece of the alfalfa biomass is surrounded by the silica polymer. The process of biomass immobilization is believed to be entrapment, and the exposed biomass as seen in Figure 4 should be responsible for the metal binding activity.

Control experiments were conducted with the silica matrix material alone and resulted in no metal binding. Figure 5 indicates the amount of metal that was bound to the




FIGURE 4. ENVIRONMENTAL SCANNING ELECTRON MICROGRAPH OF AFRICAN ALFALFA SHOOT BIOMASS. THE ALFALFA WAS EMBEDDED INSIDE A POLYSILICATE MATRIX MATERIAL USED TO SUPPORT THE BIOMASS IN COLUMNS.

column of immobilized alfalfa biomass after 120 bed volumes of 0.1 mM metal solution containing various levels of calcium and magnesium. One column was used for each heavy metal ion studied. After passing 10 bed volumes of 0.1 M HCl to recover any of the bound metal ion, the same columns were conditioned to pH 5.0 and again run with the increased concentration of hard cations. As indicated in the batch experiments, small decreases in binding was observed for copper(II) and lead(II). Significant reductions in binding were observed for cadmium(II), chromium(III), nickel(II), and zinc(II). The same trend was observed for all the binding of all the metals; as the hard cation concentration increased, the binding of heavy metals from solution slightly decreased. Nickel(II), cadmium(II), and zinc(II) showed a dramatic drop in binding upon hard cation exposure, but zinc(II) seemed to stabilize afterward. This may be




FIGURE 5. EFFECTS OF VARIOUS CALCIUM AND MAGNESIUM CONCENTRATIONS ON METAL BINDING BY SILICA-IMMOBILIZED AFRICAN ALFALFA SHOOTS UNDER FLOW CONDITIONS. CALCIUM AND MAGNESIUM CONCENTRATIONS PASSED WITH 0.1 mM METAL ION SOLUTIONS:

■ 0.0 mM ▨ 0.1 mM ▩ 10 mM ■ 100 mM

explained by the biorecovery rates for all the metals studied, as seen in Table 1. The first percent recovery rate for zinc was lower than the rest, suggesting that some of the bound zinc was not removed from the column by exposure to 0.1 M HCl, therefore hindering some of the binding. The concentration of calcium and magnesium may have also reduced the recovery of nickel(II) from the biomass. In addition, calcium and magnesium concentrations seemed to enhance the recovery of chromium(III) ions from the immobilized African alfalfa shoots. This occurrence may help in determining a better method for recovery of the strongly-bound chromium(III) from the alfalfa biomass.

CONCLUSIONS

These studies have shown that alfalfa has the potential to work like a “biological” mixed-bed-ion-exchange resin to bind heavy metals from aqueous solutions and recover these ions in a reusable form. Like ion-

exchange resins, the silica-immobilized alfalfa can be recycled, but unlike ion-exchange resins, they can be made through inexpensive methods that will not contribute environmental problems. In addition, the use of the silica-immobilized alfalfa as a “phytofilter” will allow for selective removal and recovery of various heavy metal ions, even under hard water conditions which would foul typical ion-exchange type systems. We have shown that high concentrations of hard cations do not seem to greatly reduce the binding of lead(II) and copper(II). Even with calcium and magnesium concentrations at 10,000 times that of the contaminant metals studied, the overall binding ability of silica-immobilized African alfalfa shoots was only reduced by about 30%. This innovative technology has the potential to be used for the cleanup of contaminated waters through environmentally-friendly methods.

Further studies are being performed in our laboratories to determine the chemical functional groups responsible for various metal binding by the alfalfa biomass. We would also like to conduct *in situ* experiments with actual heavy metal-contaminated waters.

TABLE 1. DESORPTION OF BOUND METAL WITH 0.1 M HCl.

	% Metal Recovered for the mM Ca & Mg Concentrations Added			
	0.0 mM	0.1 mM	10 mM	0.1 M
Cd(II)	84	85	82	70
Cr(III)	10	18	22	42
Cu(II)	90	95	100	92
Ni(II)	80	65	47	50
Pb(II)	90	100	96	93
Zn(II)	72	92	87	89

ACKNOWLEDGMENTS

The authors acknowledge financial support from the University of Texas at El Paso’s Center for Environmental Resource Management (CERM) through funding from the Office of Exploratory Research of the U.S. Environmental Protection Agency (cooperative agreement CR-819849-01-4) and the Office of Naval Research for funding of the ESEM (grant # N00014-95-1-1301). Dr. Gardea-Torresdey also acknowledges the support of NIH (grant # GM 08012-25).

REFERENCES

1. A. Alegria, R. Barberá, and R. Farré, Influence of environmental contamination on Cd, Co, Cr, Cu, Ni, Pb and Zn content of edible vegetables: Safety and nutritional aspects, *Journal of Micronutrient Analysis*, 8 (1990) 91-104.
2. W.F. Chamberlain and J.A. Miller, Barium in forage plants and in the manure of cattle treated with barium boluses, *J. Agric. Food Chem.*, 30 (1982) 463-465.
3. G. Micera and A. Dessì, Chromium adsorption by plant roots and formation of long-lived Cr(V) species: An ecological hazard?, *Journal of Inorganic Biochemistry*, 34 (1988) 157-166.
4. P.H. Kansenen and J. Venetvaara, Comparison of biological collectors of airborne heavy metals near ferrochrome and steel works, *Water, Air, and Soil Pollution*, 60 (1991) 337-359.
5. D.E. Kimbrough, Off-site forensic determination of airborne elemental emissions by multi-media analysis: A case study at two secondary lead

- smelters, *Environmental Science & Technology*, 29 (1995) 2217-2221.
6. M. Bergers and T. Harris, Lead and cadmium speciation in smelter-contaminated soil, In: L.E. Erickson, D.L. Tillison, S.C. Grant, and J.P. McDonald (Eds.), *Proceedings of the 10th Annual Conference on Hazardous Waste Research*, Kansas State Univ., Manhattan, KS, 1995, p. 56.
 7. E. Pip, Cadmium, copper, and lead in soils and garden produce near a metal smelter, Manitoba, *Bulletin Environmental Contamination and Toxicology*, 46 (1991) 790-796.
 8. C. Countryman, Arsenic and old lakes, *The Texas Observer*, Feb. 1995, pp. 6-12.
 9. National Research Council, *In Situ Bioremediation: When Does it Work?*, National Academy Press, Washington D.C., 1993, pp. 2-14.
 10. R.R. Chianelli, Bioremediation: Helping nature's microbial scavengers, In: P. Day and R. Catlow (Eds.), *The Candle Revisited, Essays on Science and Technology*, Oxford University Press, 1994, pp. 105-126.
 11. R.M. Atlas, Bioremediation, *Chem. Eng. News*, 3 (1995) 32.
 12. S. Silver, Exploiting heavy metal resistance systems in bioremediation, *Res. Microbiol.*, 145 (1994) 39-81.
 13. L.J. DeFilippi and F.S. Lupton, Bioremediation of chromium (VI) using anaerobic sulfate reducing bacteria, *American Chemical Society Division of Industrial and Engineering Chemistry*, 21-23 (1992) 117-120.
 14. J.L. Schnoor, L.A. Licht, S.C. McCutcheon, N.L. Wolfe, and L.H. Carrieria, Phytoremediation: An emerging technology for contaminated sites, In: L.E. Erickson, D.L. Tillison, S.C. Grant, and J.P. McDonald (Eds.), *Proceedings of the 10th Annual Conference on Hazardous Waste Research*, Kansas State Univ., Manhattan, KS, 1995, p. 221.
 15. W.F. Mueller, G.W. Bedell, S. Shojaee, and P.J. Jackson, Bioremediation of TNT wastes by higher plants, In: L.E. Erickson, D.L. Tillison, S.C. Grant, and J.P. McDonald (Eds.), *Proceedings of the 10th Annual Conference on Hazardous Waste Research*, Kansas State Univ., Manhattan, KS, 1995, p. 222.
 16. W.H.O. Ernst, J.A.C. Verkleij, and H. Schat, Metal tolerance in plants, *Acta Botanica Neerlandica*, Sept. (1992) 229-248.
 17. P.B.A. Nanda Kumar, V. Dushenkov, H. Motto, and I. Raskin, Phytoextraction: The use of plants to remove heavy metals from soils, *Environ. Sci. Technol.*, 29 (1995) 1232-1238.
 18. T. Adler, Botanical cleanup crews, *Science News*, 150 (1996) 42-23.
 19. Q. Xue and H.C. Harrison, Effect of soil zinc, pH, and cultivar on cadmium uptake in leaf lettuce (*Lactuca sativa* L. var. *crispa*), *Commun. Soil Sci. Plant Anal.*, 22 (1991) 975-991.
 20. R.P. Narwal, M. Singh, and J.P. Singh, Effect of nickel enriched sewage water on the accumulation of nickel and other heavy metals in corn, *J. Indian Soc. Soil Sci.*, 39 (1991) 123-128.

21. M. Vojtechova and S. Leblova, Uptake of lead and cadmium by maize seedling and the effect of heavy metals on the activity of phosphoenolpyruvate carboxylase isolated from maize, *Biologia Plantarum.*, 33 (1991) 386-394.
22. S. Sanglimsuwan, N. Yoshida, T. Moringa, and Y. Murooka, Resistance to and uptake of heavy metals in mushrooms, *J. of Fermentation and Bioengineering*, 75 (1993) 112-114.
23. R. Carrillo G. and L.J. Cajuste, Heavy metals in soils and alfalfa (*Medicago sativa* L.) irrigated with three sources of wastewater, *J. Environ. Sci. Health*, A27 (1992) 1771-1783.
24. V.C. Baligar, T.A. Campell, and R.J. Wright, Differential responses of alfalfa clones to aluminum-toxic acid soil, *J. of Plant Nutrition.*, 16 (1993) 219-233.
25. L.J. Cajuste, R. Carrillo G., E. Cota G., and R.J. Laird, The distribution of metals from wastewater in the Mexican valley of Mezquital, *Water, Air, and Soil Pollution.*, 57-58 (1991) 763-771.
26. P.W.G. Sale, D.I. Couper, P.L. Cachia, and P.J. Larkin, Tolerance to manganese toxicity among cultivar of lucerne (*Medicago sativa* L.), *Genetic Aspects of Plant Mineral Nutrition* (1993) 45-52.
27. V. Dushenkov, P.B.A. Nonda Kumar, H. Motto, and I. Raskin, Rhizofiltration: The use of plants to remove heavy metals from aqueous streams, *Environ. Sci. Technol.*, 29 (1995) 1239-1245.
28. C.M. Cooney, Sunflowers remove radionuclides from water in ongoing phytoremediation field tests, *Environ. Sci. Technol.*, 30:5 (1996) 194A.
29. C.D. Scott, Removal of dissolved metals by plant tissue, *Biotechnol. and Bioeng.*, 39 (1992) 1064-1068.
30. G.J. Ramelow, L. Liu, C. Himel, D. Fralick, Y. Zhao, and C. Tong, The analysis of dissolved metals in natural water preconcentration on biosorbents of immobilized lichen and seaweed biomass in silica, *Intern. J. Anal. Chem.*, 53 (1993) 219-232.
31. U.S. Ramelow, C.N. Guidry, and S.D. Fisk, A kinetic study of metal ion binding by biomass immobilized in polymers, *J. of Hazardous Materials*, 46 (1996) 37-55.
32. Huei-Yang D. KE and G.D. Rayson, Luminescence study of UO_2^{2+} binding to immobilized *Datura innoxia* biomaterial, *Applied Spectroscopy*, 47 (1993) 44-51.
33. B. Volesky and I. Prasetyo, Cadmium removal in a biosorption column, *Biotechnol. and Bioeng.*, 43 (1994) 1010-1015.
34. J.R. Lujan, D.W. Darnall, P.C. Stark, G.D. Rayson, and J.L. Gardea-Torresdey, Metal ion binding by algae and higher plant tissues: A phenomenological study of solution pH dependence, *Solvent Extraction and Ion Exchange*, 12:4 (1994) 803-816.
35. J.L. Gardea-Torresdey, K.J. Tiemann, J.H. Gonzalez, J.A. Henning, and M.S. Townsend, Ability of silica-immobilized *Medicago sativa* (alfalfa) to remove copper ions from solution, *J. of Hazardous Materials*, 48 (1996) 181-190.
36. J.L. Gardea-Torresdey, K.J. Tiemann, J.H. Gonzalez, I. Cano-Aguilera, J.A. Henning, and M.S. Townsend, Ability of *Medicago sativa* (alfalfa) to remove

nickel ions from aqueous solution, J. of Hazardous Materials (1996) in press.

37. J.L. Gardea-Torresdey, K.J. Tiemann, J.H. Gonzalez, J.A. Henning, and M.S. Townsend, Removal of copper ions from solution by silica-immobilized *Medicago sativa* (alfalfa), In: L.E. Erickson, D.L. Tillison, S.C. Grant, and J.P. McDonald (Eds.), Proceedings of the 10th Annual Conference on Hazardous Waste Research, Kansas State Univ., Manhattan, KS, 1995, pp. 209-217.
38. A.E. Martell and R.M. Smith, Critical Stability Constants: Other Organic Ligands, vol. 3, Plenum Press, NY and London, 1977.